

# Interstitial Cystitis/Painful Bladder Syndrome

THE UROGYNÆCOLOGICAL SOCIETY OF AUSTRALASIA

## What is interstitial cystitis/painful bladder syndrome?

Interstitial cystitis/painful bladder syndrome (IC/PBS) is a chronic condition (symptoms persist for more than 6 weeks over 3 months) that causes pain or discomfort attributed to the bladder. It is associated with a sudden desire to pass urine (urgency) as well as having to urinate very often (frequency).

Women are affected more often than men. Some women find that their symptoms are worse around the time of menstruation, when they are more stressed than usual, or when eating certain foods (triggers). Some of the symptoms are listed below:

- Urgency – during the day or night
- Frequency – sometimes up to 20x per day, as well as multiple voids overnight
- Pressure/pain around the bladder area – this gets worse as your bladder fills up and improves when you empty your bladder
- Your bladder may not hold a lot of urine
- Pain with intercourse.

## What are the causes of IC/PBS?

While the underlying cause of IC/PBS is not well understood, it is believed that a defect in the protective layer of the bladder lining results in inflammation of the underlying bladder tissue. This causes pain and irritation. Other changes include an increase in histamine (due to inflammation) and an increase in certain types of pain nerve cells in the bladder wall. There is some evidence to suggest that an autoimmune response, where the body forms antibodies against certain types of cells, also contributes to the condition.



Fig 1: Normal Bladder




Fig 2: Bladder with IC (petechiae)





Fig 3: Bladder with IC (ulcer)



# Interstitial Cystitis/Painful Bladder Syndrome

THE UROGYNÆCOLOGICAL SOCIETY OF AUSTRALASIA

## How do we treat IC?

Because we do not know the exact cause of IC, there is no one treatment that works for everyone. You and your gynaecologist may need to try several different therapies to find one that improves your symptoms.


A first-line approach is often physiotherapy and behavioural modification. This includes passing urine at specific times (timed voiding), bladder training and managing your fluid intake. It is also beneficial to identify specific 'triggers' and try to reduce these. Foods and drinks that are most bothersome include:

- Alcohol
- Coffee
- Citrus (juice/fruit)
- Cranberry juice
- Chilli peppers
- Carbonated (fizzy) drinks
- Spicy foods
- Tomatoes

As part of the cystoscopy, your gynaecologist may offer bladder distension. This means stretching the bladder wall while you are under anaesthetic. We don't know why this improves symptoms, but approximately one-third of women will get relief from with this procedure.

Medications for IC can be divided into oral medications (tablets) and instillations (solutions placed into the bladder). Medications that are available in New Zealand differ from those available in Australia, so be sure to check with your gynaecologist. Tablets that have been used include:

- Anti-inflammatories (such as ibuprofen)
- Medications that relieve neuropathic (nerve-related) pain, such as gabapentin and amitriptyline
- Anti-histaminics (such as cimetidine)
- Anticholinergics (such as oxybutynin, solifenacin or darifenacin) that relieve the urgency and frequency associated with IC
- Elmiron (pentosan polysulfate) passes into the bladder via the kidneys and coats the defects in the bladder wall. Elmiron is not funded on the PBS and you need to take it for at least 4 months to know if it will improve your symptoms





## Interstitial Cystitis/Painful Bladder Syndrome

THE UROGYNÆCOLOGICAL SOCIETY OF AUSTRALASIA

Instillations include DMSO (dimethyl sulphoxide) and hyaluronic acid. These are placed into the bladder with the help of a small catheter. The solution is left in the bladder for a maximum of 60 minutes (as long as you can tolerate it) and is then passed into the toilet. Women have one instillation per week for 6–8 weeks (this may vary between gynaecologists). Botox injections to the bladder have been shown to have a similar effect as bladder instillations, but are not currently funded for treatment of IC/PBS. Very rarely, surgery to bypass the bladder may be discussed – but this is an absolute last resort.

### More information

There are several websites where you can find useful information and IC groups in your area. Joining a support group on social media may also be helpful.

[www.ugsa.com.au](http://www.ugsa.com.au) – Information sheets on *Bladder Retraining* and *Pelvic Floor Muscle Training*

<http://www.painful-bladder.org>,

<http://www.ichelp.org/wp-content/uploads/2015/07/food-list.pdf>

(a helpful tool to evaluate food and drink 'triggers')

**This statement has been developed by the Urogynaecological Society of Australasia (UGSA).**

Disclaimer: This information is intended to provide general advice to practitioners. This information should not be relied on as a substitute for proper assessment with respect to the particular circumstances of each case and the needs of any patient. This document reflects emerging clinical and scientific advances as of the date issued and is subject to change. The document has been prepared having regard to general circumstances.

